Gary M. Annunziata, D.O., F.A.C.P / Anh T. Duong, M.D Split Dose.

NuLYTELY/ TriLyte/GoLYTELY Preparation for Colonoscopy

Your	Colonoscopy has been scheduled for	, please proceed to the:
	Mirage Endoscopy Center	
	Dolores Hope Building, First Floor R	egistration
- .		
The c	day before procedure-	
*** Yo	ou have been given a prescription OR a prescription has been	called in to your pharmacy for
NuLY	TELY/TriLyte/GoLYTELY gallon laxative and four Dulcolax La	xative tablets. We recommend that you
obtail	n either a tube of Destin ointment or A&D ointment that you ma	ay use to help with the irritation and or
tende	erness you may experience with having multiple bowel moveme	ents. ***
1)) In the morning prepare your NuLYTELY/TriLyte/GoLYTELY solution according to the instructions	
	the package and refrigerate.	
2)	Have clear liquids for breakfast, lunch and dinner. *** IT IS IM	IPORTANT THAT YOU DRINK WATER
	ALL DAY TO MAINTAIN HYDRATION. *****	
3)	3:00-4:00p.m. swallow all (4) four Dulcolax tablets with water	. The first bowel movement should occur
	in approximately 1-6 hours after this.	
4)	5:00-6:00p.m. begin drinking your NuLYTELY/TriLyte/GoLYT	ELY solution, one 8-ounce glass every
20-30 minutes until you have consumed eight glasses. THE CONTAINER WILL		CONTAINER WILL ONLY BE HALF
	FULL; PLEASE REFRIGERATE THE REST OF THE SOLUT	TION FOR THE MORNING OF
	PROCEDURE.	
5)	A watery bowel movement should occur in approximately one	e (1) hour after drinking the solution. You
	may experience some abdominal bloating, distention, and/or	nausea before bowels start to move.
	morning of the procedure:	
1) Six (6) hours prior to your procedure you will drink the remainder of the solution, one 8-ounce		•
	20-30 minutes until you have consumed the remainder of the	e solution.
2)	You may continue with clear liquids up until 4 hours prior to	your procedure
3)	A friend or relative must accompany you to drive and/or assi	st you home.